

03/04/98  
3545 U.S. PAT. & TM. OFF.

ERIDAN NEIMARK  
ROGER L BROWDY

ANNE M KORNBAU  
NORMAN J LATKER  
NICK BROMER\*  
(\*PA BAR ONLY)

OF COUNSEL  
IVER P COOPER

BROWDY AND NEIMARK, PLLC

ATTORNEYS AT LAW  
PATENT AND TRADEMARK CAUSES

SUITE 300  
419 SEVENTH STREET N W  
WASHINGTON, D. C. 20004-2299  
TELEPHONE (202)-628-5197

March 4, 1998

TELECOPIER FACSIMILE  
(202) 737-3528  
(202) 393-1012

E-MAIL  
BrwdyNmrk@digizen.net

SENIOR COUNSEL  
ALVIN BROWDY

PATENT AGENT  
ALLEN C YUN, PH D

Hon. Assistant Commissioner for Patents  
Box Patent Appln  
Washington, D.C. 20231

RE: New Patent Application in U.S.  
Applicant(s): Hajime AGA et al.  
Title: REDUCTION INHIBITORY AGENT FOR  
ACTIVE-OXYGEN ELIMINATING ACTIVITY  
Atty's Docket: AGA=6

Sir:

Attached herewith is the above-identified application for Letters Patent including:

- ☒ Specification (50 pages), claims (3 pages) and abstract (1 page)
- ☐ \_\_\_\_\_ Sheets Drawings (Figures 1- )
  - ☐ Formal ☐ Informal
- ☒ Declaration and Power of Attorney (2 pages)
  - ☒ Newly executed ☐ Copy from prior application no.
- ☐ Preliminary Amendment
- ☐ Supplemental Preliminary Amendment
- ☒ Information Disclosure Statement with 1449 and 8 references
- ☐ A verified statement to establish small entity status under 37 CFR §1.9 and 37 CFR §1.27 ( page(s))
- ☒ A check in the amount of \$ 922.00 (check no. 18010) to cover:
  - ☒ The filing fee calculated as follows:

CLAIMS AS FILED				
FOR	NUMBER FILED	NUMBER EXTRA	RATE	BASIC FEE \$ 790.00
TOTAL CLAIMS	26 - 20=	6	x 22	\$ 132.00
INDEPENDENT CLAIMS	2 - 3=	0	x 82	0
<input type="checkbox"/> Multiple Dependent Claim Presented			x270	
<input type="checkbox"/> Reduction of 1/2 for small entity				- \$
			TOTAL FILING FEE	\$ 922.00

- ☐ Any additional fee required by the filing of an enclosed preliminary or supplemental preliminary amendment has been calculated as shown below:

	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NO. PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE	CALCULATION
TOTAL	*	MINUS **	=	x \$ 22.00	\$
INDEP	*	MINUS ***	=	x \$ 82.00	\$
[ ] Multiple Dependent Claim Presented				x \$270.00	\$
Total of Above Calculations =					\$
Reduction by 1/2 for filing by small entity					- \$
Total Additional Fee =					\$


[ ] \_\_\_\_\_  
 [X] Return Receipt Postcard (in duplicate)

The following statements are applicable:

- [X] The benefit under 35 USC §119 is claimed of the filing date of: Japanese Application No. 063987/1997 in Japan on 04 March 1997, and Japanese Application No. 017647/1998 in Japan on 14 January 1998. A certified copy of said priority document [ ] is attached [ ] was filed in progenitor case \_\_\_\_\_ on \_\_\_\_\_. Application No. \_\_\_\_\_ in \_\_\_\_\_ on \_\_\_\_\_. A certified copy of said priority document [ ] is attached [ ] was filed in progenitor case \_\_\_\_\_ on \_\_\_\_\_.
- [ ] The present application is a [ ] Continuation [ ] Divisional [ ] Continuation-in-part of prior application No. \_\_\_\_\_.
- [ ] Incorporation By Reference. The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied herewith, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
- [ ] A signed statement deleting inventor(s) named in the prior application is attached.
- [ ] Amend the specification by inserting before the first line the sentence: --This is a \_\_ continuation \_\_ division of copending parent application Serial No. filed \_\_\_\_\_.--
- [ ] Certain documents were previously cited or submitted to the Patent and Trademark Office in the following prior application \_\_\_\_\_, which is relied upon under 35 USC §120. Applicants identify these documents by attaching hereto a form PTO-1449 listing these documents, and request that they be considered and made of record in accordance with 37 CFR §1.98(d). Per Section 1.98(d), copies of these documents need not be filed in this application.
- [ ] A verified statement claiming small entity status is enclosed in progenitor application no. \_\_\_\_\_, filed \_\_\_\_\_. Status is still proper and desired.
- [ ] The undersigned attorney of record hereby revokes the powers of attorney of:

- [ ] The undersigned attorney of record hereby appoints associate power of attorney, to prosecute this application and to transact all business in the Patent and Trademark Office in connection therewith to:
- [X] The Commissioner is hereby authorized to charge payment of the following additional fees associated with this communication or credit any overpayments to Deposit Account No. 02-4035:
- [X] Any additional filing fees required under 37 CFR §1.16.
  - [X] Any patent application processing fees under 37 CFR §1.17.
- [X] The Commissioner is hereby authorized to charge payment of the following fees, based on any paper filed during the pendency of this application or any CPA thereof, to effect any amendment, petition, or other action requested in said paper or credit any overpayments to Deposit Account No. 02-4035:
- [X] Any patent application processing fees under 37 CFR §1.17.
  - [ ] The issue fee set in 37 CFR §1.18 at or before mailing the Notice of Allowance, pursuant to 37 CFR §1.311(b).
  - [X] Any filing fees under 37 CFR §1.16 for presentation of extra claims.
  - [X] If a paper is untimely filed in this or any CPA thereof by Applicant(s), the Commissioner is hereby petitioned under 37 CFR §1.136(a) for the minimum extension of time required to make said paper timely. In the event a petition for extension of time is made under the provisions of this paragraph, the Commissioner is hereby requested to charge any fee required under 37 CFR §1.17 to Deposit Account 02-4035.
- [X] The Commissioner is hereby authorized to credit any overpayment of fees accompanying this paper to Deposit Account No. 02-4035.

Respectfully submitted,  
BROWDY AND NEIMARK, P.L.L.C.

By:   
Sheridan Neimark  
Registration No. 20,520

SN:bcs

f:\user3\98mar\aga6.ntr

REDUCTION INHIBITORY AGENT FOR ACTIVE-OXYGEN  
ELIMINATING ACTIVITY

Background of the Invention

Field of the Invention

The present invention relates to a reduction inhibitory agent for active-oxygen eliminating activity, uses of the same, and a method for inhibiting the reduction of the active-oxygen eliminating activity, and more particularly to a reduction inhibitory agent for active-oxygen eliminating activity, which comprises trehalose as an effective ingredient, uses of the same as compositions obtainable by incorporating the inhibitory agent into plant edible substances and/or plant antioxidants, and a method for inhibiting the reduction of active-oxygen eliminating activity of plant substances.

Description of the Prior Art

It is well known that plant edible substances such as vegetables, mushrooms, and lavers are important for living bodies as sources of functional ingredients such as vitamins, minerals, and edible fibers, and are essential food materials for the growth of living bodies. Recently, these edible plants have become to be more focused on their active-oxygen eliminating activity that relates deeply to the maintenance and promotion of health, the prevention of aging and geriatric diseases, and the inhibition of carcinogenesis. Mechanism for the cause of aging of living bodies and related diseases thereof such as cancers, arteriosclerosis, liver cirrhosis, myocardial

infarction, cerebral apoplexy, cataract, Parkinson disease, rheumatism, and Alzheimer's dementia as incurable diseases are still remained unknown in many aspects. However, the mechanism of such incurable diseases has come to be considered; active-oxygens such as superoxide which is an oxygen having unpaired electron and relatively-high reactivity, and its derivatives including hydroxyradical and hydrogen peroxide oxidize, oxidize intracellular target molecules such as membrane lipids, proteins, and DNAs to induce oxygen-related defective and to cause aging of living bodies and related diseases thereof.

Enzymes capable of eliminating active-oxygen such as superoxide dismutase (EC 1.15.1.1) and catalase (EC 1.11.1.6), and antioxidants such as L-ascorbic acid and  $\alpha$ -tocopherol exist in living cells, and the concentration of intracellular active-oxygen is kept generally to an extremely-low level. However, it is considered that irradiation of relatively-large amount of ultraviolet rays, radiations, and magnetic waves; excessive physical exercises; strong mental stresses; and aging would form active-oxygen in an amount that can not be eliminated by active-oxygen-eliminating activity of living bodies, resulting in an accumulation of compounds oxidized by the excessive amount of active-oxygen and causing the aforesaid oxygen-related defective. To improve the problem, for example, Japanese Patent Kokai Nos. 168,435/93 and 143,466/96 proposed methods which comprise supplementing to living bodies active-oxygen eliminating activity, which is easy to shortage, to promote the maintenance and promotion of health of living bodies by using a relatively-high active-oxygen eliminating activity of edible

plants. It was confirmed that even if such edible plants were used, processings such as juice extracting, extracting, heating, and drying, and successive storage conditions for the plants may lower the inherent active-oxygen eliminating activity or even extinguish the activity completely. It has been expected to establish a novel method for inhibiting the reduction of active-oxygen eliminating activity without causing serious side effects in living bodies.

### Summary of the Invention

The object of the present invention is to provide a novel reduction inhibitory agent for active-oxygen eliminating activity, a composition obtainable by incorporating the agent into plant edible substances and/or plant antioxidants to inhibit their active-oxygen eliminating activity, and a method for inhibiting the reduction of active-oxygen eliminating activity of the plant edible substance and/or the plant antioxidants.

To attain the above objects, the present inventors studied for the utilization of saccharides, and continued studying; they examined the influence of reduction inhibitory effect on active-oxygen eliminating activity of plant substances with active-oxygen eliminating activity such as plant edible substances and plant antioxidants by incorporating saccharides thereunto.

As a result, they found that trehalose exerts a more effective activity than other saccharides and inhibits strongly

the reduction of plant active-oxygen eliminating activity, and accomplished this invention.

The first object of the present invention is to provide a reduction inhibitory agent for active-oxygen eliminating activity which comprises trehalose as an effective ingredient; the second object of the present invention is to provide a composition obtainable by incorporating the agent into plant edible substances and/or plant antioxidants with active-oxygen eliminating activity to inhibit the reduction of their active-oxygen eliminating activity; and the third object of the present invention is to provide a method for inhibiting the reduction of plant active-oxygen eliminating activity, which comprises incorporating the agent into plant substances with active-oxygen eliminating activity.

#### Detailed Description of the Invention

The trehalose suitably used as an effective ingredient in the present reduction inhibitory agent for active-oxygen eliminating activity is  $\alpha,\alpha$ -trehalose, and any trehaloses can be used in the present invention independently of their origins and sources, as long as they inhibit the reduction of active-oxygen eliminating activity. For example, trehaloses prepared from yeasts as disclosed in Japanese Patent Kokai No. 246,097/95, from maltose as disclosed in Japanese Patent Kokai Nos. 216,695/83 and 170,977/95, and from starch as disclosed in Japanese Patent Kokai Nos. 143,876/95 and 213,283/95 can be arbitrarily used in the present invention. Any types of

0504336 "030498  
064030" 9843050

trehalose in the form of a syrup, crystalline powder of massecuite, crystalline hydrate, crystalline anhydride, or amorphous solid can be arbitrarily used. Commercially-available high-purity crystalline trehalose hydrate and crystalline trehalose anhydride can be arbitrarily used. The trehalose content in the present reduction inhibitory agent for active-oxygen eliminating activity can be selected from those which can allow to exert a desired reduction inhibitory activity on the active-oxygen eliminating activity of plant edible substances, etc.; usually, at least about 20 w/w % (throughout the present specification, the wording "w/w %" will be abbreviated as "%", unless specified otherwise) to the agent, on a dry solid basis (d.s.b.), and preferably at least 50%, d.s.b. The present reduction inhibitory agent can be suitably incorporated into compositions containing plant edible substances and/or plant antioxidants in an amount of, usually, at least one percent, preferably, at least about 5 percent, and more preferably at least 20%, d.s.b.

The wording "incorporating" as referred to in the present invention means "coexisting"; in the case of incorporating the present reduction inhibitory agent into plant substances with active-oxygen eliminating activity, any coexisting methods can be acceptable as long as they can allow to impart satisfactory reduction inhibitory activity for plant active-oxygen eliminating activity independently of the incorporation methods and relative proportions. The present agent can be preferably incorporated into plant edible substances and/or plant antioxidants by allowing the agent to



contact with the substances and/or the antioxidants in aqueous systems as homogeneity as possible. For example, in the case of using plant edible substances, etc., in a juicy form such as a liquid or suspension form, the present agent in the form of a powder, crystal, or solid can be incorporated into the plant edible substances as homogeneity as possible by mixing with and dissolving in the substances. While in the case of using plant edible substances and plant antioxidants in a solid form, they are treated with water to give a juicy form such as a liquid or suspension form, and incorporated with the present agent by either employing the above method or providing successively a syrup of the present agent, and either dispersing, dissolving or suspending the plant edible substances and antioxidants in the syrup to allow to contact them with the agent as homogeneity as possible and to incorporate the agent into the substances and antioxidants. In the case of using plant edible substances such as disruptant of raw plant-tissues, the present agent can be incorporated into the plant edible substances either by sprinkling over the substances and mixing them to dissolve the present agent in the substances, or by soaking the plant edible substances in the present agent in a syrup form.

The plant substances with active-oxygen eliminating activity usable in the present invention include edible plants *per se*; disrupted, minced, pulverized, dried, and extracted products of edible parts of the edible plants; and/or plant antioxidants. Examples of the edible plants suitably used in the present invention are edible rootcrops such as carrots, lotus roots, onions, burdocks, Japanese radishes, taros, and

yams; leaf vegetables such as lettuces, chicories, Chinese cabbages, cabbages, kales, *moroheiya* or *Corchorus olitorius*, *Angelica keiskei*, spinach, Malabar nightshades, *komatsuna* (a kind of Chinese cabbage), *nozawana* (a kind of turnip), *Chrysanthemum coronarium*, *chingensai* (a chinese vegetable), and turnips; fruit vegetables such as *okura* or gumbos, cauliflowers, broccolis, eggplants, tomatos, cucumbers, pumpkins, zucchini, green peppers, field peas, and kidney beans; vegetables such as alfalfa, bean sprouts including those of soy beans and mung beans; mushrooms such as *shiitake* mushrooms, winter mushrooms or *Celtis sinensis*, and *shimeji* mushrooms; seaweeds such as *hijiki* seaweed or *Hijikia fusiformis*, *wakame* seaweed or *Undaria pinnatifida*, and tangs; citrus fruits such as lemons, citrons, and *sudachi* (a kind of citrus fruit), shaddocks, and kumquats; fruits such as bananas, kiwifruits, strawberries, hawthorns, blueberries, grapes, peaches, apples, pears, chestnuts; herbs such as garlics, gingers, *wasabis* (a Japanese horseradish), mustards, parsleys, Chinese parsleys, beefsteak plants, leeks, Welsh onions, celeries, dropworts, cressons, Guinea peppers, Japanese peppers, peppers, rosemaries, and mints; medicinal herbs and crude drugs such as mugworts, common plantains, bad-smelling perennial plants of the family *Saururaceae*, *Cassia obtusifolia*, Japanese green gentians, aloes, licorices, turmeric, Japanese indigo plants, *Pfaffia*, loquat leaves, field horsetails, pine needles, bamboo leaves, umes or Japanese apricots, tea leaves, barley leaves, wheat leaves, oats leaves, rye leaves, buckwheat leaves, ginkgo leaves, Chinese gutta

percha leaves, *oobanasarusuberi* (a plant of the family *Lythrum*), *Aspalathus linearis*, and *Gymnema sylvestre*; and others including edible parts of nuts, seeds, and germs such as adlays, buckwheats, sesames, rices, wheats, corns, broad beans, soybeans, peanuts, walnuts, pine seeds, and other seeds and germs.

One or more of the following relatively-low-molecular-weight plant antioxidants can be suitably selected and used as the present plant antioxidants: Plant enzymes having active-oxygen eliminating activity such as superoxide dismutase, catalase, and superoxide; plant pigments such as rutin,  $\alpha$ -glucosyl rutin, hesperidin,  $\alpha$ -glucosyl hesperidin, naringin,  $\alpha$ -glucosyl naringin, chlorophyll, carotenoid, and anthocyanin; plant polyphenols such as gallic acid, catechin,  $\alpha$ -glucosyl catechin, and tannic acid; and plant vitamins such as  $\alpha$ -tocopherol, L-ascorbic acid, and riboflavin. In the present invention, the relatively-low-molecular-weight plant antioxidants can be contained in the edible plants, or added supplementarily to the contents during the processing of the edible parts of edible plants. The plant antioxidants, which exist separately to the edible plants, can be used in the present invention. The plant antioxidants should not be restricted to those which are extracted from plants and include those which are artificially synthesized and produced.

Explaining more concretely, processes for producing the present composition, which the reduction of the active-oxygen eliminating activity is satisfactorily inhibited, are as follows: Edible parts of edible plants are, for example, (i)

disrupted or cut by mixers, juicers, or pulverizes into pastes or suspensions, into which the present agent is incorporated by adding thereto and dissolving therein; or (ii) extracted with solvents such as hot water and alcohols, and filtered, if necessary, further concentrated into liquids or pastes, followed by incorporating the present agent into the liquids or pastes by mixing to dissolve therein. The compositions thus obtained are usually syrupy, pasty, or juicy disrupted products which have a relatively-high moisture content and require a relatively-low storage temperature for a satisfactory shelf-life. The compositions can be dried and/or pulverized, or dried and/or pulverized after sterilization by heating to obtain powdery or small pieces of solid compositions. Conventional drying and pulverizing methods can be arbitrarily used in the present invention; the compositions of syrupy, pasty, or juicy disrupted products can be (i) successively dried *in vacuo*, air-dried, drum-dried, pulverized, and sieved; (ii) successively spray-dried, subjected to fluidized-drying, and sieved; or (iii) successively mixed with anhydrous saccharides such as maltose or trehalose anhydride to effect dehydration, dried, and sieved into powdery products. If necessary, the powdery products can be arbitrarily granulated, tabletted, and capsulated into appropriate shapes and forms in use. The solid compositions, processed up to such a high-level, will acquire more improved activity for inhibiting the reduction of active-oxygen eliminating activity, storage stability, and handleability. The solid compositions containing edible plant substances have features that they inhibit effectively the deterioration of

plant pigments, contained in the material edible plants, such as chlorophylls, carotenoids, anthocyanin, and flavonoids; retain satisfactorily the inherent color of the pigments; and mask unfavorable tastes such as bitterness and astringency.

The present composition, containing trehalose and a plant edible substance or plant antioxidant and having inhibitory activity for the reduction of active-oxygen eliminating activity, can be further mixed with one or more appropriate substances in the form of a solid or liquid such as essential minerals, edible fibers, saccharides for promoting the growth of bifid bacteria, vitamins, biologically active substances, and preservatives.

The essential minerals usable in the present include, for example, calcium, magnesium, phosphorus, iron, copper, zinc, and cobalt, and these minerals are used in an adequate amount. The edible fibers satisfactorily usable in the present invention are, for example, pectin, alginic acid, carrageenan, gum arabic, glucomannan, cyclodextrin, and pullulan. It was revealed that among the edible fibers, particularly pullulan exerts a strong reduction-inhibitory-activity for plant active-oxygen eliminating activity when used in combination with trehalose as the effective ingredient according to the present invention. Based on this, pullulan can be most satisfactorily used together with trehalose in the present agent and used in the production of the present compositions.

Examples of the saccharides used for promoting the growth of bifid bacteria in the present invention are lactosucrose, fructooligosaccharide, galactooligosaccharide, and

isomaltooligosaccharide which can be arbitrarily used on demands.

The vitamins satisfactorily usable in the present invention include, for example, water-soluble vitamins such as thiamin, riboflavin, L-ascorbic acid, rutin, hesperidin, naringin, niacin, pyridoxine, cyanocobalamine, and derivatives thereof; and lipid-soluble vitamins such as vitamin A, vitamin D,  $\alpha$ -tocopherol, vitamin K, and derivatives thereof.

One or more of biological active substances such as hormones, antibiotics, cytokines, and propolis can be arbitrarily used on demands. Adequate amounts of ethanol, acetic acid, lactic acid, and salt can be arbitrarily used on demands as preservatives in the present invention.

To improve the quality of the present compositions and other compositions containing the present compositions, adequate coloring agents, flavoring agents, taste-imparting agents, stabilizers, fillers, adjuvants, and excipients can be arbitrarily used.

The compositions thus obtained have at least five units/g composition, d.s.b., and more preferably, at least 10 units/g composition, d.s.b., of active-oxygen eliminating activity as determined by the nitroblue tetrazolium (NBT) test, and they can be arbitrarily and easily administered to subjects via appropriate routes of oral, intubational, percutaneous, and permucosal administrations to impart active-oxygen eliminating activity with lesser deterioration of their quality. Intake of the present composition containing plant edible substances enriches plant functional ingredients such as vitamins,

minerals, and edible fibers inherent to the plants, and enriches living bodies with active-oxygen eliminating activity. Thus the present composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and to inhibit carcinogenesis; and arbitrarily used as food products, cosmetics, pharmaceuticals, and their materials and processing intermediates.

To enrich nutritional value and improve qualities, taste, and taste-preference, the present composition can be arbitrarily used for health food products and other food products in general, for example, seasonings such as "*furikake*" (a seasoned fish meal), sauces, ketchup, "*yakiniku-no-tare*" (a sauce for Japanese grilled meat), curry roux, and instant soup mixes; Japanese and Western cakes such as "*mochi*" (a rice paste), "*dango*" (a ball of rice paste), candies, chewing gums, backed confectioneries, snack confectioneries, waffles, sponge cakes, buns, and bakeries; frozen desserts such as ice creams, sherbets, and candies; pastes such as fruit pastes, fruit sources, peanut pastes, and raw jams; pickles and pickled products; meat products such as hams and sausages; products of fish meats such as "*kamaboko*" (a steamed fish paste), "*chikuwa*" (a kind of fish paste), "*hanpen*" (a fish cake), and sausages; "*chinmi*" (a delicacy); "*tsukudani*" (a food boiled down in soy); daily dishes; beverages such as vegetable juices, fruit juices, and carbonated beverages; rice products such as rice gruels, porridges of rice and vegetables, and cooked rice with seasonings, meats, and vegetables; noodles such as iced noodles,

macaronis, and pastas; food products such as mix for pudding, premixes for hot cakes, instant juices, instant soups, and frozen foods; and their materials and processing intermediates. The present composition can be arbitrarily used in cosmetics such as skin-beautifying agents, packs, creams, shampoos, hair rinses, hair tonics, bath salts, enzymic agents, and tooth pastes in the form of a liquid, paste, powder, or granule; and used as a material or a processing intermediate for those products. Furthermore, the present composition can be used to treat and prevent aging-related diseases in the form of an oral and/or intubation nutrient or therapeutic agent such as a collunarium, nebula, digestive agent, stomachic, enzymic agent, ointment, cataplasma, and their materials and processing intermediates.

The present composition consists substantially of food materials and can be used with less fear of causing toxicity. Usually it can be suitably used on a dry solid basis at a dose of about 0.1 to 1,000 g/adult/day, preferably, about 0.2 to 500 g/adult/day, and more preferably, about 1 to 100 g/adult/day when administered to subjects orally. The present composition can be applied to subjects intubationally, percutaneously, and permucosally in accordance with the dose of the oral administration.

The following experiments describe the present invention in detail:

#### Experiment 1

Influence of saccharides on reduction inhibitory activity for active-oxygen eliminating activity



864030" 93343050

Fresh raw carrots were disrupted by a mixer, and 10% of either of saccharides, d.s.b., was added to the mixture and dissolved therein. The solution was dried under reduced pressures at 40°C for two days, dried *in vacuo* at the same temperature for 24 hours, and pulverized by a mixer into a powdery carrot composition. The saccharides used were commercialized reagents of glucose, mannitol, sorbitol, maltose, sucrose, trehalose, and pullulan. An about 100 g of the composition was placed and sealed in a 500-ml polystyrene container, and stored at 40°C for seven days. The active-oxygen eliminating activity of the composition before and after the storage were compared. The assay for the active-oxygen eliminating activity followed the nitroblue tetrazolium test by Toshio IMANARI et al in *Igakuno Ayumi*, Vol. 101, pp. 496-497 (1977); the level of superoxide, formed in a xanthine-xanthine oxidase system, was assayed by quantifying calorimetrically the content of formazan formed by reducing NBT. As a control, refined water was used as a test solution. One unit activity of active-oxygen eliminating activity is defined as that inhibits 50% formazan formation as compared with the control. Table 1 shows the composition of the powdery compositions, and the active-oxygen eliminating activity per gram of either of the compositions before and after the storage. As evident from Table 1, among the saccharides tested the composition with trehalose showed the highest residual percentage for active-oxygen eliminating activity, revealing that trehalose exerts a strong effect on the reduction inhibitory activity for active-oxygen eliminating activity. The reduction inhibitory activity

of trehalose was strongly higher than that of glucose as the constituent saccharide for trehalose, and that of maltose as a disaccharide consisting of two glucose units similar to trehalose. Thus trehalose is suitably used as an effective ingredient for a reduction inhibitory agent for active-oxygen eliminating activity. Next to trehalose, pullulan showed a relatively-high reduction inhibitory activity for active-oxygen eliminating activity.

Table 1

Saccharide	Composition of powdery composition (%)		Saccharide content (d.s.b.) (%)	Activity (unit/g)		Percentage of residual activity (%)	
	Carrot Saccharide Moisture			Before storage	After storage		
None	93.4	0.0	6.6	0.0	870	100	11
Glucose	48.8	46.4	4.8	48.7	580	170	29
Mannitol	49.8	47.5	2.7	48.8	520	59	11
Sorbitol	48.2	46.0	5.8	48.8	560	26	5
Maltose	48.1	45.8	6.1	48.8	550	180	33
Sucrose	48.2	46.0	5.8	48.8	660	180	27
Trehalose	47.9	45.7	6.4	48.8	580	380	66
Pullulan	47.7	45.5	6.8	48.8	710	360	51

## Experiment 2

### Influence of trehalose concentration on reduction inhibitory activity for active-oxygen eliminating activity

Similarly as in Experiment 1, 0-20% of trehalose, d.s.b., was mixed with and dissolved in a disrupted carrot. The resulting each mixture was dried in vacuo at 40°C for 24 hours, and pulverized into a powdery carrot composition. Similarly as in Experiment 1, each powdery composition was placed and sealed in a polystyrene container, and stored at 40°C for seven days for evaluating the reduction inhibitory activity for active-oxygen eliminating activity. Table 2 shows the active-oxygen eliminating activity of one gram of each powdery composition before and after the storage. Based on the data from Table 2, at least of about one percent, preferably, at least of about five percent, and more preferably, at least of about 20% of trehalose, d.s.b., exerted a desired effective reducing inhibitory activity for active-oxygen eliminating activity.

Table 2

Percentage (%) of trehalose added to carrot	Composition of powdery composition (%)		Trehalose content (d.s.b.) (%)	Activity (unit/g)		Percentage of residual activity (%)	
	Carrot	Trehalose Moisture		Before storage	After storage		
0.0	95.3	0.0	4.7	0.0	940	330	35
0.1	94.3	1.0	4.7	1.0	930	400	43
0.5	90.4	4.7	4.9	4.9	750	460	61
1.0	85.9	8.9	5.2	9.4	860	480	56
2.5	75.3	19.6	5.1	20.7	700	460	66
5.0	62.3	32.4	5.3	34.2	600	370	62
10.0	46.1	48.1	5.8	51.1	400	330	83
20.0	30.3	63.1	6.6	67.6	380	240	63

### Experiment 3

#### Influence of trehalose on reducing inhibitory activity for active-oxygen eliminating activity of vegetables

Vegetables were treated similarly as in Experiment 1, and 10% trehalose, d.s.b., to the disrupted vegetables. The resulting each mixture was dried *in vacuo* at 45°C for 20 hours, and pulverized into a powdery vegetable composition. Similarly as in Experiment 1, each compositions was respectively placed in a container and stored at 40°C for six days. The active-oxygen eliminating activity of one gram of each composition before and after the storage was compared with that of the composition prepared without trehalose. Table 3 shows the composition of the powdery compositions, and their active-oxygen eliminating activity. Reduction inhibitory activity for active-oxygen eliminating activity by incorporation of trehalose was observed in vegetables such as carrot, onion, Japanese radish, cabbage, spinach, cucumber, and pumpkin.

Table 3

Vegetable (moisture (%))	Composition of powdery composition (%)		Trehalose content (d.s.b.) (%)	Activity (unit/g)		Percentage of residual activity (%)
	Carrot	Saccharide Moisture		Before storage	After storage	
Carrot (90.1)	94.5	0.0	5.5	1200	370	31
	46.2	46.6	7.2	570	460	81
Onion (89.3)	93.2	0.0	6.8	440	210	48
	47.5	44.3	8.2	300	230	77
Japanese radish (94.0)	92.7	0.0	7.3	1100	630	57
	35.0	58.3	6.7	530	450	85
Cabbage (93.8)	94.8	0.0	5.2	4100	2500	63
	35.7	57.5	6.8	1600	1400	88
Spinach (94.1)	93.2	0.0	6.8	3500	1300	36
	34.4	58.4	7.2	1200	890	74
Eggplant (93.7)	91.5	0.0	8.5	38000	31000	82
	34.8	55.3	9.9	13000	11000	85
Cucumber (95.0)	92.7	0.0	7.3	4600	1300	30
	30.3	60.5	9.2	1900	1100	58
Pumpkin (84.0)	93.2	0.0	6.8	1600	610	38
	56.2	35.2	8.6	880	460	52

#### Experiment 4

#### Reduction inhibitory activity for active-oxygen eliminating activity by trehalose with respect to superoxide dismutase and antioxidants

In one milliliter of a 45 w/v % trehalose solution was dissolved five micrograms of superoxide dismutase from a horseradish commercialized by Sigma Chemical Co., St. Louise, USA, 10 mg  $\alpha$ -glucosyl hesperidin commercialized by Hayashibara Biochemical Laboratories, Inc., Okayama, Japan, 10 mg  $\alpha$ -glucosyl rutin commercialized by Hayashibara Biochemical Laboratories, Inc., Okayama, Japan, 250  $\mu$ g gallic acid commercialized by Wako Pure Chemical Industries, Ltd., Tokyo, Japan, or 50  $\mu$ g catechin to obtain a liquid composition. Each of the compositions was placed in a glass test tube, sealed with a rubber cap, and stored under light-shielded conditions. As a control, water was used in place of the trehalose solution. Table 4 shows the active-oxygen eliminating activity before and after the storage. As shown in Table 4, it was confirmed that trehalose exerts an effective reducing inhibitory activity for active-oxygen eliminating activity of plant antioxidants such as superoxide dismutase,  $\alpha$ -glucosyl hesperidin,  $\alpha$ -glucosyl rutin, gallic acid, and catechin. Mere mixing of a powdery crystalline trehalose hydrate and either of the powdery antioxidants into a powdery mixture, almost no change of the active-oxygen eliminating activity was observed when subjected to storage test similarly as above. Based on the results, it was revealed that a reducing inhibitory agent for active-oxygen eliminating activity, which comprises trehalose as an effective ingredient, could scarcely



exert the desired activity only when mixed in a powder form with powdery substances having active-oxygen eliminating activity, but exert the desired activity when trehalose in a melted state is allowed to contact with and incorporated into the substances, or when trehalose is incorporated into the substances in an aqueous system.

864080" SEE 4E060

Table 4

Content of antioxidant	Storage condition	Trehalose concentration (%)	Activity (unit/mg antioxidant)		Residual activity (%)
			Before storage	After storage	
Superoxide dismutase (5 µg/ml)	60° C 17 hours	0 45	10,000 11,000	4,600 9,400	46 85
α-Glucosyl hesperidin (10 mg/ml)	80° C 5 days	0 45	3.8 3.6	1.0 2.4	26 67
α-Glucosyl rutin (10 mg/ml)	80° C 5 days	0 45	25 26	18 22	72 85
Gallic acid (250 µg/ml)	80° C 5 days	0 45	590 570	390 470	66 82
Catechin (50 µg/ml)	25° C 20 hours	0 45	470 380	2.2 310	0 82

## Experiment 5

### Reduction inhibitory effect for active-oxygen eliminating activity on herb extracts

Ten liters of water was added to 500 g of a dried specimen of plantain, *Plantago asiatica*, commercialized by Daido Seiyaku Co., Ltd., Tokyo, Japan, and the mixture was infused for 90 min, followed by collecting a supernatant of the infused solution by a basket-type centrifuge. The supernatant was boiled down to obtain two kilograms of an about 5 w/v % solution. Two hundred and forty grams of trehalose and 10 g of pullulan, d.s.b., were mixed with and dissolved in one kilogram of the above concentrate, and the solution was spray dried by "SD-1", a portable spray-dryer commercialized by Tokyo Rika Mfg. Co., Ltd., Tokyo, Japan, Tokyo, Japan, into a composition containing an extract of *Plantago asiatica*. As a control, using dextrin with DE (dextrose equivalent) of about 11 in place of the trehalose and pullulan, a powdery composition similar to the above composition was prepared. Similarly as in Experiment 1, these powdery compositions were placed in polystyrene containers, sealed, and stored at 40°C for seven days for examining the reduction inhibitory effect for active-oxygen eliminating activity. Both the compositions had an activity of about 5,000 units/g composition before the storage. On seven days after the storage, the composition with trehalose and pullulan had an activity of about 4,300 units/g composition, while the composition with dextrin had an activity of about 3,100 units/g composition. The data shows that the use of trehalose and pullulan exerts a higher reduction inhibitory

effect for active-oxygen eliminating activity than in the case of using dextrin.

The following Examples describe the present invention; Examples A describe the present reducing inhibitory agent for active-oxygen eliminating activity, and Examples B describe the present composition where the reduction of active-oxygen eliminating activity is satisfactorily inhibited:

Example A-1

Reduction inhibitory agent for active-oxygen eliminating activity

Corn starch was prepared into an about 33% starch suspension which was then mixed with 0.1% calcium carbonate, adjusted to Ph 6.5, mixed with 0.3% per g starch of "TERMAMYL", an  $\alpha$ -amylase specimen commercialized by Novo Industri A/S Copenhagen, Denmark, enzymatically reacted at 95°C for 15 min, autoclaved at 120°C, and promptly cooled to obtain a liquefied solution with DE of about four. Thereafter, the liquefied solution was admixed with four units/g starch, d.s.b., of respective non-reducing saccharide-forming enzyme and trehalose-releasing enzyme, prepared by culturing a microorganism of the genus *Rhizobium* by the method in Japanese Patent Kokai No. 213,283/95, 500 units/g starch of isoamylase, and five units/g starch of cyclomaltodextrin glucanotransferase, adjusted to Ph 6.2, and enzymatically reacted at 40°C for 48 hours. The reaction mixture was heated to inactivate the remaining enzymes, then admixed with 10 units/g substrate of glucoamylase and enzymatically reacted at pH 5.0 and 50°C for 10 hours. The reaction mixture thus obtained contained about 86% trehalose,



00043303049  
SECRET  
maltose, and about 6% glucose. The reaction mixture was in a conventional manner purified by decoloring and desalting, concentrated *in vacuo* into an about 75% syrup which was then placed in a crystallizer, admixed with about one percent of respective seed crystals of crystalline trehalose hydrate and crystalline maltose hydrate, and cooled to obtain a massecuite with a crystallization degree of about 25%. The massecuite was spray dried and aged to obtain a powdery crystalline saccharide in a yield of about 92% to the material maltose, d.s.b. The powdery saccharide is a stable powdery saccharide, which has a crystallization degree of about 60% and contains about 66% trehalose, about 28% maltose, and about six % glucose, and it can be suitably used as a reduction inhibitory agent for active-oxygen eliminating activity and arbitrarily used to inhibit the reduction of active-oxygen eliminating activity of plant edible substances.

#### Example A-3

#### Reduction inhibitory agent for active-oxygen eliminating activity

Potato starch was prepared into a 10% starch suspension which was then subjected to the action of  $\alpha$ -amylase to obtain a liquefied solution. To the liquefied solution were added three units/g starch, d.s.b., of a non-reducing saccharide-forming enzyme disclosed in Japanese Patent Kokai No. 213,283/95, five units/g starch, d.s.b., of a trehalose-releasing enzyme, 1,000 units/g starch, d.s.b., of an isoamylase, and one unit/g starch, d.s.b., of a maltotetraose-forming amylase, and the mixture was enzymatically reacted at

pH 6.0 and 40°C for 48 hours. The reaction mixture was heated to inactivate the remaining enzyme, and in a conventional manner purified by decoloring and desalting and concentrated *in vacuo* to obtain a syrup with a moisture content of about 30% and DE of about 15 in a yield of about 90%, d.s.b. The product is a stable syrup, which contains, on a dry solid basis, about 50% trehalose along with other saccharides derived from starch, can be suitably used as a reduction inhibitory agent for active-oxygen eliminating activity, and arbitrarily used to inhibit the reduction of active-oxygen eliminating activity of plant edible substances.

Example A-4

Reduction inhibitory agent for active-oxygen  
eliminating activity

One hundred parts by weight of a high-purity crystalline trehalose hydrate, obtained by the method in Example A-1, was mixed to homogeneity with one part by weight of pullulan to produce a solid reduction inhibitory agent for active-oxygen eliminating activity. The product can be arbitrarily used in a reduction inhibitory agent for active-oxygen eliminating activity of plant edible substances.

Example A-5

Reduction inhibitory agent for active-oxygen  
eliminating activity

In 100 hundred parts by weight of a trehalose syrup, obtained by the method in Example A-3, was dissolved by mixing to homogeneity 0.5 part by weight of pullulan, and 0.5 part by weight of a readily-water-soluble cyclodextrin into a syrupy

reduction inhibitory agent for active-oxygen eliminating activity. The product can be arbitrarily used in a reduction inhibitory agent for active-oxygen eliminating activity of plant edible substances.

#### Example B-1

##### Composition containing spinach

Fresh spinach was pulverized by a mixer, and in 10 parts by weight of the resulting disruptant were dissolved by mixing one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-4, and 0.1 part by weight of  $\alpha$ -glucosyl rutin. The resulting mixture was heated at 100°C for 10 min, dried with air heated to 40°C for two hours, dried *in vacuo* at 40°C for 16 hours, and powdered by a pulverizer to obtain a composition containing spinach having an active-oxygen eliminating activity of about 1,200 units/g composition.

The composition retained satisfactorily the inherent color of spinach and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the spinach, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-2



#### Composition containing cabbage

Cabbage was cut, branched, sliced by a cutter into fine strips in about 5 mm wide. Ten parts by weight of the fine strips was mixed with one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-3, and 0.05 part by weight of  $\alpha$ -glucosyl rutin, and the mixture was allowed to stand at ambient temperature for two hours, then dried by air heated to 40°C for two hours, dried *in vacuo* at 40°C for 16 hours to obtain a composition containing fine strips of cabbage having an active-oxygen eliminating activity of about 1,300 units/g composition. The composition retained satisfactorily the inherent color of the cabbage and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the cabbage, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-3

#### Composition containing eggplant

A fresh eggplant was subjected to a slicer, and the sliced eggplant was washed with water. One part by weight of a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-1, was mixed with

and dissolved in 10 parts by weight of the sliced eggplant, and the resulting mixture was dried by air heated to 70°C for two days to obtain a composition containing sliced eggplant having an active-oxygen eliminating activity of about 11,000 units/g composition. The composition retained the inherent color of the eggplant's skin and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the eggplant, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-4

##### Composition containing carrot

A fresh carrot was branched and sliced by a slicer. One part by weight of a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-4, was added to 10 parts by weight of the sliced carrot and dissolved therein. The mixture was dried with air heated to 50°C for 16 hours to obtain a composition containing sliced carrots having an active-oxygen eliminating activity of about 450 units/g composition. The composition satisfactorily retained the color inherent to the carrot, stimulated your appetite, and had a relatively-low hygroscopicity and a satisfactory shelf-life. Intake of the composition enriches



or processing intermediate.

Example B-6

Composition containing *shiitake* mushroom

A dried *shiitake* mushroom was swelled out in water, heated at 100°C for 15 min, and disrupted by a cutter. In 100 parts by weight of the disrupted mixture was dissolved by mixing 0.1 part by weight of a tea extract and 10 parts by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-3, and the resulting mixture was dried at 50°C for eight hours, dried *in vacuo* at 40°C, and pulverized to obtain a composition containing a powdery *shiitake* mushroom having an active-oxygen eliminating activity of about 560 units/g composition. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to *Hizikia fusiforme*, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-7

Composition containing citron

Citron rind was disrupted by a cutter, and five parts by weight of the disrupted rind was mixed with one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-1, to

004033-000498  
dissolve the agent therein. After preliminary freezing, the resulting mixture was freeze-dried for three days, and disrupted to obtain a composition containing a powdery citron having an active-oxygen eliminating activity of about 12 units/g composition. The composition retains satisfactorily the color and flavor inherent to the citron. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the citron, and imparts the active-oxygen eliminating activity to living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-8

##### Composition containing ginger

A fresh ginger was disrupted by a mixer. In five parts by weight of the disrupted mixture was dissolved by mixing one part by weight of an reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-2. After preliminary freezing, the resulting mixture was freeze-dried for three days, and disrupted by a crusher to obtain a composition containing a powdery ginger having an active-oxygen eliminating activity of about 120 units/g composition. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the ginger, and imparts the active-

oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-9

Composition containing aojiso (a beefsteak plant)

Aojiso leaves were branched and disrupted by a mixer. In five parts by weight of the disrupted mixture was dissolved by mixing one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-4, and 0.5 part by weight of  $\alpha$ -glucosyl rutin, then mixed with 49 parts by weight of crystalline trehalose anhydride as a desiccant. The resulting mixture was allowed to stand at ambient temperature for one day and disrupted by a crusher to obtain a composition containing a powdery aojiso having an oxygen eliminating activity of about 230 units/g composition. The composition satisfactorily retained the color, flavor and taste of the aojiso, and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the aojiso, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the

composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-10

##### Composition containing mugwort

According to a conventional manner, mugwort was placed in a boiling water with an adequate amount of salt to remove harshness, dehydrated softly and disrupted by a crusher. Five parts by weight of the disrupted mixture was mixed with 0.05 part by weight of  $\alpha$ -glucosyl rutin and 1.3 parts by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-5. The resulting mixture was dried by air heated to 50°C for four hours, dried *in vacuo* at 40°C for 16 hours, and disrupted to obtain a composition containing a powdery mugwort having an active-oxygen eliminating activity of about 780 units/g composition. The composition satisfactorily retained the color, flavor and taste of the mugwort, and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to mugwort, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-11

##### Composition containing dokudami (a bad-smelling

perennial plant of the family Saururaceae

A half part by weight of a dried *dokudami* was mixed with 10 parts by weight of water, and heated to boiling for 90 min. A supernatant of the infused solution was collected by a basket-type centrifuge and boiled down to obtain four parts by weight of an about five w/v % solution. Two parts by weight of the concentrate was mixed with 0.5 part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-4, and the resulting solution was spray-dried in a conventional manner to obtain a composition containing a powdery *dokudami* extract having an active-oxygen eliminating activity of about 1,400 units/g composition. The composition had a lesser unsatisfactory smell and taste inherent to the *dokudami* and could be more easily swallowable than conventional ones. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the *dokudami*, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-12

##### Composition containing aloe

A fresh aloe was disrupted by a mixer, and 10 parts by weight of the disrupted aloe was mixed with 0.1 part by



weight of a tea extract and one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-1. The resulting mixture was dried by air heated to 70°C for two days and disrupted by a crusher to obtain a composition containing a powdery aloe having an active-oxygen eliminating activity of 5,800 units/g composition. The composition had a lesser astringency of the aloe and was more easily swallowable than conventional ones. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the aloe, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-13

##### Composition containing Chinese cabbage preserved with seasonings

A Chinese cabbage preserved with seasonings was subjected to a cutter. To 10 parts by weight of the cut product was added and dissolved therein one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-1, and the mixture was dried with air heated to 55°C for 16 hours to obtain a composition containing small pieces of Chinese cabbage, preserved with seasonings, having an active-oxygen eliminating activity of about 5,400 units/g composition. The product

retained the color tint of the Chinese cabbage. When you tasted the product in your mouth, the satisfactory taste and flavor spread throughout your mouth and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the Chinese cabbage, and augments the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate. The composition can be also used as a food product such as *furikake*, premix for *onigiri* (a rice ball), and soup for Chinese noodles, and as a health food, cosmetic, pharmaceutical, or their material or intermediate.

#### Example B-14

##### Composition of nozawana (a kind of turnip)

A pickled nozawana was subjected to a cutter. To 10 parts by weight of the cut product was added and dissolved therein one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-1. The mixture was dried with air heated to 55°C for 16 hours to obtain a composition containing small pieces of pickled nozawana having an active-oxygen eliminating activity of about 2,600 units/g composition. The product retained the color tint of the pickled nozawana. When you tasted the product in your mouth, the satisfactory taste and flavor spread

throughout your mouth and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the pickled nozawana, and augments the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

#### Example B-15

##### Composition of Korean pickles of Chinese cabbage

A Korean pickles of Chinese cabbage was subjected to a cuter. To 10 parts by weight of the cut product was added and dissolved therein one part by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-4. The mixture was dried with air heated to 55°C for 16 hours to obtain a composition containing small pieces of pickled nozawana having an active-oxygen eliminating activity of about 1,600 units/g composition. The product retained the color tint of the Korean pickles of Chinese cabbage. When you tasted the product in your mouth, the satisfactory taste and flavor spread throughout your mouth and stimulated your appetite. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the Korean pickles of Chinese cabbage, and augments the active-oxygen eliminating activity to the living bodies. The composition can be

864030-9E4E050

satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-16

Composition containing soy bean

Soy beans were soaked in five-fold volumes of water overnight, and drained. Ten parts by weight of the resulting soy beans was placed in a pun, mixed with 35 parts by weight of a reduction inhibitory agent for active-oxygen eliminating activity obtained by the method in Example A-1, and 35 parts by weight of water. The mixture was boiled over a mild fire up to give a concentration of about 70% as determined by a brix apparatus and to obtain boiled beans. The beans were placed in a basket and dried with air heated to 40°C overnight to obtain a composition containing soy beans having an active-oxygen eliminating activity of about 25 units/g composition. The product retained the color tint of the boiled soy beans. When you tasted the product in your mouth, the satisfactory taste and flavor spread throughout your mouth and stimulated your appetite. In the product, there exists crystalline trehalose hydrate within the soy beans' tissues, and the hydrate well retained the tissues. The product was a solid food having satisfactory taste and biting property. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the soy beans, and augments the active-oxygen eliminating activity

to the living bodies. Thus the composition can be satisfactorily used to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-17

Composition containing apple

Fresh apples were peeled and subjected to a slicer to obtain sliced apples, about 5 mm in thickness, which were then soaked for 30 min in an about 75% solution prepared by dissolving a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-1, in water by heating at 80°C, placed in a basket, and dried at 35°C overnight to obtain a composition containing apple having an active-oxygen eliminating activity of about 50 units/g composition. The product retained the color tint of the apple. When you tasted the product in your mouth, the satisfactory taste and flavor spread throughout your mouth and stimulated your appetite. In the product, there exists crystalline trehalose hydrate within the apple's tissues, and the hydrate well retained the tissues. The product was a solid food having satisfactory taste and biting property. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the soy beans, and augments the active-oxygen eliminating activity to the living bodies. Thus the composition can be satisfactorily used to maintain and promote your health, prevent

aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-18

Composition containing carrot

Fresh carrots were subjected to a slicer, and five parts by weight of the sliced carrots was placed in an about 73% syrup which was mildly boiling at about 105°C and which had been prepared by dissolving by heating 140 parts by weight of a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-1, in 60 parts by weight of water. The resulting mixture was continued concentrating by boiling up to give a temperature of about 110°C for 20 min and up to be dehydrated by heating into a fried-like product. Thereafter, the product was placed in a basket, dried by air heated to 50°C for five hours to obtain a composition containing sliced carrots having an active-oxygen eliminating activity of about 200 units/g composition. The composition was a fried-like product of carrot, to which crystalline trehalose hydrate adhered, and it retained the inherent color of the carrot and stimulated your appetite. The product had a relatively-low hygroscopicity, satisfactorily shelf-life, and desired handleability. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the carrot, and imparts the active-oxygen eliminating activity to the living bodies. The composition can be satisfactorily used to maintain and

promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and prevent carcinogenesis. Thus the composition can be arbitrarily used as a food product, cosmetic, pharmaceutical, or their material or processing intermediate.

Example B-19

Composition containing apple

Fresh apples were peeled and subjected to a slicer into sliced apples, about 5 mm in thickness. Four parts by weight of the sliced apples was then placed in an about 80% syrup heated to about 110°C, which had been prepared by dissolving by heating in water 140 parts by weight of a reduction inhibitory agent for active-oxygen eliminating activity, obtained by the method in Example A-1, and continued concentrating by boiling up to give a temperature of about 114°C for about 10 min and up to be dehydrated by heating into a fried-like product. Thereafter, the product was placed in a basket, dried by air heated to 35°C overnight to obtain a composition containing sliced apples having an active-oxygen eliminating activity of about 30 units/g composition. The composition was a fried-like product of apple, to which crystalline trehalose hydrate adhered, and it retained the inherent color of the apple and stimulated your appetite. The product had a relatively-low hygroscopicity, satisfactorily shelf-life, and desired handleability. Intake of the composition enriches living bodies with functional ingredients such as vitamins, minerals, and edible fibers inherent to the apple, and imparts the active-oxygen eliminating activity to the





manner shaped and packaged into a *dango* having an active-oxygen eliminating activity of over 12 units/g content. The product is a *yomogi-dango*, i.e., a rice paste with mugwort, having a satisfactory color tint of *yomogi*, flavor, taste, and biting property.

#### Example B-22

##### Nutrition

A composition consisting of the following ingredients was prepared; 20 parts by weight of crystalline  $\alpha$ -maltose, 1.1 parts by weight of glycine, 0.18 part by weight of sodium glutamate, 1.2 parts by weight of salt, one part by weight of citric acid, 0.4 part by weight of calcium lactate, 0.3 part by weight of a composition containing spinach obtained by the method in Example B-1, 0.01 part by weight of thiamine, and 0.01 part by weight of riboflavin. Twenty-four grams aliquots of the composition were injected into small laminated aluminum bags and heat sealed to obtain a nutrition for intubation having an active oxygen eliminating activity of over 12 units/g composition. One bag of the nutrition is dissolved in about 300-500 ml water, and the solution can be arbitrarily administered to subjects orally or to the subjects' nasal cavities, stomachs, and intestines as a parenteral liquid-supplemental-nutrition.

#### Example B-23

##### Cosmetic cream

Two parts by weight of polyoxyethylene glycol monostearate, 5 parts by weight of glyceryl monostearate, self-emulsifying, 2.8 parts by weight of a composition containing

aloe obtained by the method in Example B-12, 0.2 part by weight of  $\alpha$ -glucosyl rutin, one part by weight of liquid petrolatum, 10 parts by weight of glyceryl tri-2-ethylhexanoate, and an adequate amount of an antiseptic were dissolved by heating in a conventional manner. The resulting solution was admixed with two parts by weight of L-lactic acid, five parts by weight of 1,3-butylene glycol and 66 parts by weight of refined water, and the mixture thus obtained was emulsified by a homogenizer and admixed with an adequate amount of a flavor under stirring conditions to obtain a cosmetic cream having an active-oxygen eliminating activity of over 46 units/g product. The product can be arbitrarily used as a therapeutic or preventive agent for sunburn skin, skin-beautifying agent, skin-whitening agent, and agent for inhibiting skin-aging-phenomena such as chloasmas, freckles, pigmentation, and wrinkles.

#### Example B-24

##### Bath salt

A bath salt having an active oxygen eliminating activity of over 220 units/g product, was prepared by mixing 26 parts by weight of refined water and adequate amounts of a coloring agent and a flavoring agent with 21 parts by weight of sodium DL-lactate, eight parts by weight of sodium pyruvate, five parts by weight of a composition containing citron obtained by the method in Example B-7, one part by weight of  $\alpha$ -glucosyl rutin, and 40 parts by weight of ethanol. The product can be suitably used as a skin-beautifying agent and skin-whitening agent by diluting 100-10,000 fold in hot water in a bathtub. Similarly as above, the product can be arbitrarily used by

diluting in water for face wash or beauty wash before use.

#### Example B-25

##### Ointment

One part by weight of sodium acetate trihydrate and four parts by weight of calcium DL-lactate were mixed to homogeneity with 10 parts by weight of glycerine, and the mixture was added to a mixture consisting of 50 parts by weight of petrolatum, 10 parts by weight of vegetable wax, five parts by weight of lanoline, 14.5 parts by weight of sesame oil, six parts by weight of a composition containing *dokudami* obtained by the method in Example B-11, and 0.5 part by weight of peppermint oil. The mixture thus obtained was further mixed to homogeneity into an ointment having an active-oxygen eliminating activity of about 54 units/g product. The product can be arbitrarily used as an antipruritic agent, skin-beautifying agent, skin-whitening agent, and agent for promoting the treatment for traumas and burns.

#### Example B-26

##### Tablet

Twenty parts by weight of ascorbic acid were mixed to homogeneity with 13 parts by weight of crystalline  $\beta$ -maltose, four parts by weight of corn starch, three parts by weight of a composition containing plantain and trehalose obtained by the method in Experiment 5 according to the present invention. The resulting mixture was tableted into a tablet using a 20R punch, 12 mm in diameter. The tablet had an oxygen eliminating activity of over 560 units/g product, and can be administered orally to subjects 1-10 tablets/adult/day. Especially, it can

be used for removing fever and maintaining/controlling stomach and intestinal conditions.

As described above, the present invention provides a reduction inhibitory agent for active-oxygen eliminating activity which comprises trehalose as an effective ingredient, a method for inhibiting the reduction of active-oxygen eliminating activity characterized in that it comprises incorporating either trehalose or the reduction inhibitory agent into products to be treated, and a composition which contains plant an edible substance and/or a plant antioxidant where the reduction of active-oxygen eliminating activity is satisfactorily inhibited by the above method. Intake of the present composition easily supplements to living bodies and enriches living bodies with and functional ingredients such as vitamins, minerals, and edible fibers of plant edible substances. Therefore the present composition contributes greatly to maintain and promote your health, prevent aging and geriatric diseases, promote the treatment of incurable diseases, and inhibit carcinogenesis. Thus the establishment of the present invention would explore a novel health resource as a fourth functional ingredient usable in the fields of processings and applications of plant edible substances, and will give a great contribution to a wide variety of fields, especially, industrial fields of food products, cosmetics, and pharmaceuticals.

While there has been described what is at present considered to be the preferred embodiments of the invention, it

will be understood the various modifications may be made therein, and it is intended to cover in the appended claims all such modifications as fall within the true spirits and scope of the invention.

SEE PAGE 50

WE CLAIMS:

1. A reduction inhibitory agent for active-oxygen eliminating activity, which comprises trehalose as an effective ingredient.

2. The inhibitory agent of claim 1, which further contains a saccharide other than said trehalose.

3. The inhibitory agent of claim 1, which contains pullulan and/or cyclodextrin.

4. The inhibitory agent of claim 1, which contains at least 20 w/w % of said trehalose, on a dry solid basis.

5. A method for inhibiting the reduction of active-oxygen-eliminating activity, which comprises incorporating the inhibitory agent of claim 1, into a plant substance with active-oxygen eliminating activity in an aqueous-system.

6. The method of claim 5, wherein said plant substance is a plant edible substance and/or a plant antioxidant.

7. The method of claim 5, wherein said plant edible substance is in the form of a sliced, disrupted, pulverized, dried, or extracted plant-edible-part.

8. The method of claim 6, wherein said plant antioxidant is a member selected from the group consisting of plant enzymes, pigments, polyphenols, and vitamins.

9. The method of claim 5, wherein at least about one w/w %, on a dry solid basis, of said trehalose as an effective ingredient is incorporated into said plant substance.

10. The method of claim 5, which contains a

sterilization step and/or a drying step.

11. A composition obtainable by incorporating said inhibitory agent of claim 1 into a plant substance with active-oxygen eliminating activity to inhibit the reduction of said activity.

12. The composition of claim 11, wherein said plant substance is a plant edible substance and/or a plant antioxidant.

13. The composition of claim 12, wherein said plant edible substance is an organ and/or tissue of an edible plant, or a sliced, disrupted, pulverized, dried, pickled, or extracted plant-edible-part.

14. The composition of claim 12, wherein said plant antioxidant is a member selected from the group consisting of plant enzymes, pigments, polyphenols, and vitamins.

15. The composition of claim 11, which contains at least about one w/w %, on a dry solid basis, of said trehalose as an effective ingredient.

16. The composition of claim 11, which contains said trehalose in an amorphous or crystalline hydrous form.

17. The composition of claim 11, which contains one or more members selected from the group consisting of essential minerals, edible fibers, growth-promoting saccharides for bifid bacteria, vitamins, and biologically active substances.

18. The composition of claim 11, which has been received a sterilization step and/or a drying step.

19. The composition of claim 11, which has active-oxygen eliminating activity of at least five units/g

composition, on a dry solid basis, as determined by the nitroblue tetrazolium test.

20. The composition of claim 11, which is in the form of a food product, cosmetic, pharmaceutical, or their material or intermediate.

21. A food product, cosmetic, or pharmaceutical, into which said composition of claim 11 is incorporated.

22. A composition into which at least about 30 w/w % of trehalose is incorporated into an edible plant tissue, on a dry solid basis.

23. The composition of claim 22, wherein said edible plant tissue is a member selected from the group consisting of an edible plant's organs and tissues, and their sliced and pickled products where trehalose is incorporated into said edible plant tissue in an amorphous or crystalline hydrous form.

24. The composition of claim 22, which has been received a sterilization and/or drying steps.

25. The composition of claim 22, which is in the form of a solid food product.

26. The composition of claim 25, wherein said solid food product is for enjoying a satisfactory biting property.



Abstract of the Disclosure

A reduction inhibitory agent for active-oxygen eliminating activity which comprises trehalose as an effective ingredient, a method for inhibiting the reduction of active-oxygen eliminating activity which comprises incorporating either trehalose or the reduction inhibitory agent into plant edible products and/or plant antioxidants, and a composition where the reduction of active-oxygen eliminating activity of the plant edible products and/or plant antioxidants is satisfactorily inhibited by the method.

05034336-030498

## Combined Declaration for Patent Application and Power of Attorney

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled (insert full title here) **REDUCTION INHIBITORY AGENT FOR ACTIVE-OXYGEN ELIMINATING ACTIVITY**

the specification of which (check one)

☒ [xx] is attached hereto;

☐ [ ] was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
USSN \_\_\_\_\_\*; or

☐ [ ] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an intentional (PCT) application, PCT/ \_\_\_\_\_; filed \_\_\_\_\_, entry requested on \_\_\_\_\_\*; national stage application received USSN \_\_\_\_\_\*; §371/§102(e) date \_\_\_\_\_\* (\*if known), and was amended on \_\_\_\_\_ (if applicable).

(include dates of amendments under PCT Art. 19 and 34 if PCT)

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119, 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>63987/1997</u>	<u>Japan</u>	<u>4th March 1997</u>	<input checked="" type="checkbox"/> [x]	<input type="checkbox"/> [ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO
<u>17647/1998</u>	<u>Japan</u>	<u>14th January 1998</u>	<input checked="" type="checkbox"/> [x]	<input type="checkbox"/> [ ]
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. § 120 of any prior U.S. non-provisional Application(s) or prior PCT application(s) designating the U.S. listed below, or under § 119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

_____ (Application Serial NO.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application Serial NO.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)

I hereby appoint the following attorneys, with full power of substitution, association, and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

SHERIDAN NEIMARK, REG. NO. 20,520 - ROGER L. BROWDY, REG. NO. 25,618 - ANNE M. KORNBAU, REG. NO. 25,884  
NORMAN J. LATKER, REG. NO. 19,963 - IVER P. COOPER, REG. NO. 28,005 - ALLEN C. YUN, REG. NO. 37,971\*  
NICK S. BROMER, REG. NO. 33,478 - \*Patent Agent

ADDRESS ALL CORRESPONDENCE TO  
**BROWDY AND NEIMARK, P.L.L.C.**  
419 Seventh Street, N.W.  
Washington, D.C. 20004

DIRECT ALL TELEPHONE CALLS TO  
**BROWDY AND NEIMARK**  
(202) 628-5197

The undersigned hereby authorizes the U.S. Attorneys or Agents named herein to accept and follow instructions from SUMA PATENT OFFICE as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorney or Agent and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents named herein will be so notified by the undersigned.

"SECRET" 060426

Title: REDUCTION INHIBITORY AGENT FOR ACTIVE-OXYGEN ELIMINATING ACTIVITY

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR Hajime AGA		INVENTOR'S SIGNATURE <i>Hajime Aga</i>	DATE Feb. 17, 1998
RESIDENCE Okayama, Japan		CITIZENSHIP Japanese	
POST OFFICE ADDRESS 880-225, Minato, Okayama-shi, Okayama, Japan			
FULL NAME OF SECOND JOINT INVENTOR Takashi SHIBUYA		INVENTOR'S SIGNATURE <i>Takashi Shibuya</i>	DATE Feb. 17, 1998
RESIDENCE Okayama, Japan		CITIZENSHIP Japanese	
POST OFFICE ADDRESS 318, Shimobara, Souja-shi, Okayama, Japan			
FULL NAME OF THIRD JOINT INVENTOR Shigeharu FUKUDA		INVENTOR'S SIGNATURE <i>Shigeharu Fukuda</i>	DATE Feb. 17, 1998
RESIDENCE Okayama, Japan		CITIZENSHIP Japanese	
POST OFFICE ADDRESS 2189, Atsu, Okayama-shi, Okayama, Japan			
FULL NAME OF FOURTH JOINT INVENTOR Toshio MIYAKE		INVENTOR'S SIGNATURE <i>Toshio Miyake</i>	DATE Feb. 17, 1998
RESIDENCE Okayama, Japan		CITIZENSHIP Japanese	
POST OFFICE ADDRESS 3-23, 1-chome, Ishima-cho, Okayama-shi, Okayama, Japan			
FULL NAME OF FIFTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SIXTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SEVENTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS.

RECEIVED 9204060